

FISHER INFORMATION IN k -RECORDS

GLENN HOFMANN^{1*} AND N. BALAKRISHNAN²

¹*Departamento de Estadística, Facultad de Ciencias Físicas y Matemáticas, Barrio Universitario, Casilla 160-C, Concepción, Chile, e-mail: glennhofmann@yahoo.com*

²*Department of Mathematics and Statistics, McMaster University, 1280 Main Street West, Hamilton, Ontario, Canada L8S 4K1, e-mail: bala@mcmaster.ca*

(Received July 9, 2002; revised February 3, 2003)

Abstract. We derive some general results on the Fisher information (FI) contained in the upper (or lower) k -record values and associated k -record times generated from an i.i.d. sample of fixed size from a continuous distribution. We apply the results to obtain the FI in both upper and lower k -record data from an exponential distribution. We propose two estimators of the exponential mean, based on the upper and lower k -record data, and discuss their small sample properties. We also consider k -record data from an inverse sampling plan, and present general formulas for the FI contained in it.

Key words and phrases: Record values, record times, k -record values, Fisher information, exponential distribution, parameter estimation, inverse sampling.

*Supported in part by Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) grant # 1020479 of Chile.