

Testing equality between several populations covariance operators

Graciela Boente 1 · Daniela Rodriguez 2 · Mariela Sued 3

Received: 14 December 2015 / Revised: 22 May 2017 / Published online: 20 August 2017 © The Institute of Statistical Mathematics, Tokyo 2017

Abstract In many situations, when dealing with several populations, equality of the covariance operators is assumed. An important issue is to study whether this assumption holds before making other inferences. In this paper, we develop a test for comparing covariance operators of several functional data samples. The proposed test is based on the Hilbert–Schmidt norm of the difference between estimated covariance operators. In particular, when dealing with two populations, the test statistic is just the squared norm of the difference between the two covariance operators. The asymptotic behaviour of the test statistic under both the null hypothesis and local alternatives is obtained. The computation of the quantiles of the null asymptotic distribution is not feasible in practice. To overcome this problem, a bootstrap procedure is considered. The performance of the test statistic for small sample sizes is illustrated through a Monte Carlo study and on a real data set.

Keywords Asymptotic distribution · Bootstrap calibration · Covariance operators · Functional data analysis · Local alternatives

Mariela Sued marielasued@gmail.com

> Graciela Boente gboente@dm.uba.ar

Daniela Rodriguez drodrig@dm.uba.ar

- ¹ Departamento de Matemáticas, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires and IMAS, CONICET, Ciudad Universitaria, Pabellón 1, 1428 Buenos Aires, Argentina
- ² Instituto de Cálculo, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires and IMAS, CONICET, Ciudad Universitaria, Pabellón 2, 1428 Buenos Aires, Argentina
- ³ Instituto de Cálculo, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires and CONICET, Ciudad Universitaria, Pabellón 2, 1428 Buenos Aires, Argentina