

Multiscale inference for a multivariate density with applications to X-ray astronomy

Konstantin Eckle¹ · Nicolai Bissantz¹ · Holger Dette¹ · Katharina Proksch² · Sabrina Einecke³

Received: 13 May 2016 / Revised: 17 November 2016 / Published online: 24 February 2017 © The Institute of Statistical Mathematics, Tokyo 2017

Abstract In this paper, we propose methods for inference of the geometric features of a multivariate density. Our approach uses multiscale tests for the monotonicity of the density at arbitrary points in arbitrary directions. In particular, a significance test for a mode at a specific point is constructed. Moreover, we develop multiscale methods for identifying regions of monotonicity and a general procedure for detecting the modes of a multivariate density. It is shown that the latter method localizes the modes with an effectively optimal rate. The theoretical results are illustrated by means of a simulation study and a data example. The new method is applied to and motivated by the determination and verification of the position of high-energy sources from X-ray observations by the Swift satellite which is important for a multiwavelength analysis of objects such as Active Galactic Nuclei.

Keywords Multiple tests · Modes · Multivariate density · X-ray astronomy

Holger Dette Holger.dette@rub.de

¹ Fakultät für Mathematik, Ruhr-Universität Bochum, 44780 Bochum, Germany

² Institut f
ür Mathematische Stochastik, Georg-August-Universit
ät G
öttingen, 37077 G
öttingen, Germany

³ Fakultät Physik, Technische Universität Dortmund, 44221 Dortmund, Germany