

# A LOWER BOUND FOR THE NUMBER OF SINGULAR SATURATED MAIN EFFECT PLANS OF AN $S^m$ FACTORIAL\*

B. L. RAKTOE AND W. T. FEDERER

(Received April 24, 1970)

## Summary

A fraction consisting of  $m(s-1)+1$  observations taken at  $m(s-1)+1$  treatment combinations of an  $s^m$  factorial with the aim to estimate the mean and the  $m(s-1)$  main effect single degree of freedom parameters is called a saturated main effect plan. If the design matrix of such a fraction is singular, then the fraction is called a singular saturated main effect plan. This paper presents a lower bound on the number of singular saturated main effect plans.

## 1. Introduction

Consider the full replicate of an  $s^m$  factorial ( $s$  is a prime or a power of a prime), then it is well known (e.g. see Kempthorne [3], that the  $s^m$  treatment combinations form the points of the finite Euclidean geometry  $EG(m, s)$  over the field  $GF(s)$  and that the  $(s^m-1)/(s-1)$  effects are 1:1 correspondence with the points of the finite projective geometry  $PG(m-1, s)$ . Each effect of the  $s^m$  factorial represents  $(s-1)$  degree of freedom, which in fact means that every point  $(x_1, x_2, \dots, x_m)$  of  $PG(m-1, s)$  represents the class  $\rho(x_1, x_2, \dots, x_m)$ , where  $\rho$  is a non-zero element of  $GF(s)$ . The elements of a class can then be taken to depict a set of  $(s-1)$  single degree of freedom parameters. If we adjoin the mean  $\mu = (A^0 B^0 \dots M^0)$  to all single degree of freedom parameters, then it is readily seen that the  $s^m$  single degree of freedom parameters are in 1:1 correspondence with the points of  $EG(m, s)$ . Hence we have explicitly that the set of treatment combinations  $\{(z_1, z_2, \dots, z_m), z_i \in GF(s)\}$  is in a 1:1 correspondence with the set of single degree of freedom parameters  $\{(A^1 B^2 \dots M^m), (z_1, z_2, \dots, z_m) \in EG(m, s)\}$ . This means that if we wish to discuss properties of both the treatment combinations and the single degree of freedom parameters we may limit ourselves to the finite Euclidean geometry  $EG(m, s)$ .

\* Research supported by N.R.C. Grant No. A07204.

If we have a full replicate of an  $s^m$  factorial, then the usual linear model tying up the observations and the single degree of freedom parameters (assuming that the factors have quantitative levels) is:

$$(1.1) \quad E[Y] = X\beta$$

where:  $Y$  is an  $s^m \times 1$  vector of observations, each component of which is taken at a treatment combination,  $X$  is an  $s^m \times s^m$  matrix such that  $X'X = \text{diagonal}(d_1, d_2, \dots, d_N)$ ,  $N = s^m$ , and  $\beta$  is an  $s^m \times 1$  vector of single degree of freedom parameters as described earlier.

DEFINITION 1.1. Define an element  $A^{z_1}B^{z_2}\dots M^{z_m}$  of  $\beta$  to be a main effect single degree of freedom parameter if the superscript  $(z_1, z_2, \dots, z_m)$  has exactly one non-zero coordinate.

DEFINITION 1.2. A plan consisting of  $m(s-1)+1$  observations to estimate the mean  $\mu$  and the  $m(s-1)$  main effect single degree of freedom parameters is termed a saturated main effect plan.

Now, if the  $[m(s-1)+1]$ -vector of observations is denoted by  $Y_1$ , then we know that the normal equations for a saturated main effect plan is:

$$(1.2) \quad X'_{11} X_{11} \hat{\beta}_1 = X'_{11} Y_1$$

where:  $X_{11}$  is an  $[m(s-1)+1] \times [m(s-1)+1]$  matrix simply read off from  $X$  of equation (1.1),  $\hat{\beta}_1$  is the least squares estimator of  $\beta$ , which is the  $[m(s-1)+1]$ -vector with  $\mu = A^0 B^0 \dots M^0$  as its first element and the rest of the elements being main effect single degree of freedom parameters.

DEFINITION 1.3. Following Banerjee and Federer [1] we define a saturated main effect plan to be singular if  $\text{rank}[X_{11}] < m(s-1)+1$  and nonsingular if  $\text{rank}[X_{11}] = m(s-1)+1$ .

The question naturally arises as to which type of plans will yield singular saturated main effect plans or equivalently which  $[m(s-1)+1]$ -subsets of  $EG(m, s)$  will give rise to singular  $X_{11}$  matrices. Also, if singular saturated main effect plans exist, then we wish to know how many of the

$$(1.3) \quad \binom{s^m}{m(s-1)+1}$$

possible plans are singular. These two questions can be considered as the general problem of singular (or nonsingular if the complementary problem is considered) saturated main effect plans.

The aim of this paper is to discuss a sub-class of singular saturated main effect plans, which lends itself to be treated geometrically and is

in agreement with the theory of confounding. This then will lead us to a lower bound on the number of singular saturated main effect plans.

## 2. Existence and enumeration of singular saturated main effect plans

Consider an incomplete block design consisting of  $s^m$  treatments in  $s^k$  blocks of  $s^{m-k}$  plots each, then it is well known (see for example Kempthorne [3]) that the construction of such a design can always be done by providing a confounding scheme such that  $(s^k-1)/(s-1)$  effects are completely confounded with the blocks. This statement of course is equivalent to providing a  $(k-1)$ -flat of  $PG(m-1, s)$ , which in turn is equivalent to the exhibition of its set of  $k$  generators.

Suppose now that  $G = \{A^{x_{11}}B^{x_{12}} \dots M^{x_{1m}}, A^{x_{21}}B^{x_{22}} \dots M^{x_{2m}}, A^{x_{k1}}B^{x_{k2}} \dots M^{x_{km}}\}$  is a set of  $k$  generators of a particular confounding scheme or equivalently of a particular  $(k-1)$ -flat of  $PG(m-1, s)$ , then the following definition will be needed in the sequel:

DEFINITION 2.1. The  $(\alpha_1, \alpha_2, \dots, \alpha_k)$ th level of a set of  $k$  generators, denoted by the symbol,

$$(2.1) \quad \{A^{x_{11}}B^{x_{12}} \dots M^{x_{1m}}, A^{x_{21}}B^{x_{22}} \dots M^{x_{2m}}, \dots, A^{x_{k1}}B^{x_{k2}} \dots M^{x_{km}}\}_{(\alpha_1, \alpha_2, \dots, \alpha_k)}$$

will be defined to depict a set of  $s^{m-k}$  treatment combinations  $\{(z_1, z_2, \dots, z_m)\}$  in  $EG(m, s)$ , satisfying the consistent and independent set of equations:

$$(2.2) \quad \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1m} \\ x_{21} & x_{22} & \dots & x_{2m} \\ \cdot & \cdot & \cdot & \cdot \\ x_{k1} & x_{k2} & \dots & x_{km} \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ \cdot \\ z_m \end{bmatrix} = \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \cdot \\ \alpha_k \end{bmatrix}$$

where  $(\alpha_1, \alpha_2, \dots, \alpha_k)$  is a given set of  $k$  elements from  $GF(s)$ . Note that the solutions to (2.2) form a  $(m-k)$ -flat of  $EG(m, s)$ . Also note, that for a fixed set of  $k$  generators of a confounding scheme we have precisely  $s^k$  fractional replicates of order  $s^{m-k}$ , since each of the  $\alpha_i$ 's can be chosen in  $s$  ways from  $GF(s)$ . A particular fractional replicate of this type can then be denoted by:

$$(2.3) \quad I = \{A^{x_{11}}B^{x_{12}} \dots M^{x_{1m}}, A^{x_{21}}B^{x_{22}} \dots M^{x_{2m}}, \dots, A^{x_{k1}}B^{x_{k2}} \dots M^{x_{km}}\}_{(\alpha_1, \alpha_2, \dots, \alpha_k)}$$

with the usual meaning that the mean is completely confounded with  $(s^k-1)/(s-1)$  effects, generated by the  $k$  generators within the braces.

DEFINITION 2.2. Define  $k$  to be the largest positive integer such

that we may select  $m(s-1)+1$  treatment combinations (i.e. a saturated main effect plan) from among the  $s^{m-k}$  combinations of the fraction (2.3). This definition of  $k$  implies the following inequalities for given  $m$  and  $s$ :

$$(2.4) \quad m(s-1)+1 \leq s^{m-k}$$

$$(2.5) \quad \text{i.e.} \quad m \leq (s^{m-k}-1)/(s-1).$$

If we denote a saturated main effect plan of  $m(s-1)+1$  treatment combinations selected from among  $s^{m-k}$  treatment combinations where  $k$  satisfies (2.4) or (2.5), by  $D$ , then the following can be easily verified:

**THEOREM 2.1.** *The number of plans of type  $D$  is given by*

$$(2.6) \quad N[m-1, k-1, s] = \phi[m-1, k-1, s] \cdot s^k \cdot \binom{s^{m-k}}{m(s-1)+1}$$

where  $\phi[m-1, k-1, s]$  is the number of  $(k-1)$ -flats in  $PG(m-1, s)$  explicitly given by (see Mann [4]):

$$(2.7) \quad \phi[m-1, k-1, s] = \left[ \prod_{i=0}^{k-1} (s^{m-i}-1) \right] / \left[ \prod_{i=0}^{k-1} (s^{k-i}-1) \right].$$

Of course, the number given by (2.6) is less than the number  $T(m, s)$  of saturated plans selected in an unrestricted manner, i.e.

$$(2.8) \quad N[m-1, k-1, s] < \binom{s^m}{m(s-1)+1} = T(m, s).$$

Now, let  $T[m, s, 0]$  denote the number of all singular saturated main effect plans, i.e. those plans from among the  $T(m, s)$  plans which lead to singular  $X_{11}$  matrices in our setting (1.2), i.e., those plans for which  $|X_{11}|=0$ . (The number  $T[m, s, 0]$  is not known for all  $s^m$  factorials and this problem and some related ones are currently under study.) Let  $N[m-1, k-1, s, 0]$  denote the total number of singular saturated main effect plans of type  $D$  as described earlier, then our intention is to determine  $N[m-1, k-1, s, 0]$  and to show that this number is a lowerbound to  $T[m, s, 0]$ . Naturally we have to discuss first the existence of singular saturated main effect plans. The following theorem establishes the existence of singular saturated main effect plans:

**THEOREM 2.2.** *If a main effect or a two-factor interaction is completely confounded with the mean, then the fraction leads to a singular saturated main effect plan.*

**PROOF.** Let  $k$  satisfy the inequality (2.4) or (2.5) and let  $m(s-1)+1$  treatment combinations be selected from the fraction

$$(2.9) \quad I = \{A^{x_{11}}B^{x_{12}} \cdots M^{x_{1m}}, \dots, A, \dots, A^{x_{k1}}B^{x_{k2}} \cdots M^{x_{km}}\}_{(a_1, a_2, \dots, a_k)}.$$

Here we have for simplicity, but without loss of generality, chosen the main effect  $A$  ( $A$  represents, as pointed out in Section 1, a set of  $(s-1)$  main effect single degree of freedom parameters). From (2.9) it follows immediately that in the design matrix  $X_{11}$  of (1.2) the columns corresponding to  $A^{u_1=1}, A^{u_2}, \dots, A^{u_{s-1}}$  will have columns of the form  $c_1 1, c_2 1, \dots, c_{s-1} 1$ , where the  $u_i$ 's are non-zero elements of  $GF(s)$  and the  $c_i$ 's are integers and finally 1 is an  $[m(s-1)+1]$ -column vector of  $+1$ 's. Hence it follows that  $\text{rank } [X_{11}] < m(s-1)+1$ , i.e.  $X_{11}$  is singular. Similarly if we select  $m(s-1)+1$  combinations from the fraction:

$$(2.10) \quad I = \{A^{x_{11}} B^{x_{12}} \dots M^{x_{1m}}, \dots, AB, \dots, A^{x_{k1}} B^{x_{k2}} \dots M^{x_{km}}\}_{(\alpha_1, \alpha_2, \dots, \alpha_k)}$$

where without loss of generality we have taken the two factor interaction  $AB$ , then by the usual group theoretic multiplication we obtain that  $A$  is completely confounded with  $B$ . This then as above immediately implies the singularity of  $X_{11}$ .

This theorem implies that the class of all singular saturated main effect plans of type  $D$  is completely characterized by the set of confounding schemes in which either a main effect or two-factor interaction is confounded, this set arising from the consideration of an incomplete block design with  $s^m$  treatments in  $s^k$  blocks of  $s^{m-k}$  plots each, where  $k$  satisfies either (2.4) or (2.5). The number  $N[m-1, k-1, s, 0]$  of such plans can be determined from the following theorem.

**THEOREM 2.3.** *Let  $k$  be the largest integer such that for given  $m$  and  $s$  the inequality (2.4) or (2.5) is satisfied and let  $L$  be the set of elements of  $PG(m-1, s)$  having exactly one or two coordinates not equal to zero, then the number  $H[m-1, k-1, s]$  of  $(k-1)$ -flats in  $PG(m-1, s)$  incident with a point of  $L$  is given by:*

$$(2.11) \quad H[m-1, k-1, s] = \phi[m-1, k-1, s] - \theta[m-1, k-1, s]$$

where

$$(2.12) \quad \theta[m-1, k-1, s] = \frac{\prod_{i=0}^{m-1} [(s^{m-k}-1) - (s-1)i]}{s^{(m-k)(m-k-1)/2} \prod_{i=0}^{m-k-1} (s^{m-k-i}-1)}.$$

**PROOF.** This theorem has been proved by Dowling [2] in a more general setting and he will publish the results shortly in a separate paper. However, note that the theorem as stated above is just a re-statement of a combinatorial confounding problem in terms of the finite projective geometry  $PG(m-1, s)$ .

From Theorem 2.2 and Theorem 2.3 we then have the following corollary:

COROLLARY 2.1. *The number  $N[m-1, k-1, s, 0]$  of singular saturated main effect plans of type D is given by:*

$$(2.13) \quad N[m-1, k-1, s, 0] = H[m-1, k-1, s] \cdot s^k \cdot \binom{s^{m-k}}{m(s-1)+1}.$$

Next, we claim the following theorem:

THEOREM 2.4.  *$N[m-1, k-1, s, 0]$  is a lowerbound to  $T[m, s, 0]$ .*

PROOF. It is sufficient to exhibit for one particular  $m$  and  $s$  a set of  $m(s-1)+1$  treatment combinations which is not of type of  $D$  and which leads to a singular saturated main effect plan. Consider the plan:  $\{(0000), (0011), (0101), (1111), (1010)\}$  for the  $2^4$  factorial. The matrix  $X_{11}$  is then

$$(2.14) \quad X_{11} = \begin{matrix} & \mu & A/2 & B/2 & C/2 & D/2 \\ \begin{pmatrix} 1 & -1 & -1 & -1 & -1 \\ 1 & -1 & -1 & 1 & 1 \\ 1 & -1 & 1 & -1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & 1 & -1 \end{pmatrix} \end{matrix}.$$

By inspection of the plan we see that no main effect nor any two factor interaction is completely confounded with the mean. Also can be verified easily that  $|X_{11}|=0$ . Finally, in some cases  $N[m-1, k-1, s, 0]$  is equal to  $T[m, s, 0]$ , e.g. in the  $2^3$  factorial. Hence we have the conclusion:

$$(2.15) \quad N[m-1, k-1, s, 0] \leq T[m, s, 0].$$

### 3. Discussion

The determination of  $T[m, s, 0]$  is not so easy and this problem belongs to a more general problem of determining the values which the determinant  $|X_{11}|$  can assume when arbitrary selections of  $m(s-1)+1$  treatment combinations are made from among  $s^m$  treatment combinations. The next problem is then to find exactly how many selections belong to a particular determinant. There are of course various ways in which these problems can be solved and methods of attacks are being studied in extenso.

## REFERENCES

- [1] Banerjee, K. S. and Federer, W. T. (1966). On estimation and construction in fractional replication, *Ann. Math. Statist.*, **37**, 1033-1039.
- [2] Dowling, Th. A. (1970). Private communication.
- [3] Kempthorne, O. (1952). *The design and analysis of experiments*, Wiley, New York.
- [4] Mann, M. B. (1949). *Analysis and design of experiments*, Dover Publications, Inc., New York.

Addendum To  
"A LOWER BOUND FOR THE NUMBER OF SINGULAR SATURATED  
MAIN EFFECT PLANS OF AN  $S^M$  FACTORIAL"

B. L. RAKTOE AND W. T. FEDERER

The following omissions should be added to the paper which appeared in the Ann. Inst. Statist. Math., 22 (1970), 519-525:

*Line 23 on page 522* should read: ...as described earlier *such that a main effect or two-factor interaction is completely confounded with the mean*, then...

*Line 15 on page 523* should read: ...implies that *a subclass of the class of all...*

*Line 7 on page 524* should read: ...not of type D of *Theorem 2.2* and...