

# Uncertainty principle and quantum Fisher information

Paolo Gibilisco · Tommaso Isola

Received: 15 January 2006 / Revised: 15 September 2006 /  
Published online: 30 December 2006  
© The Institute of Statistical Mathematics, Tokyo 2006

**Abstract** A family of inequalities, related to the uncertainty principle, has been recently proved by S. Luo, Z. Zhang, Q. Zhang, H. Kosaki, K. Yanagi, S. Furuichi and K. Kuriyama. We show that the inequalities have a geometric interpretation in terms of quantum Fisher information. Using this formulation one may naturally ask if this family of inequalities can be further extendend, for example to the *RLD* quantum Fisher information. We show that this is impossible by producing a family of counterexamples.

**Keywords** Uncertainty principle · Monotone metrics · Quantum Fisher information · Wigner–Yanase–Dyson information

## 1 Introduction

Noncommutativity in quantum probability has far-reaching consequences. One of the most important is the Heisenberg uncertainty principle

$$\text{Var}_\rho(A) \cdot \text{Var}_\rho(B) \geq \frac{1}{4} |\text{Tr}(\rho[A, B])|^2.$$

---

P. Gibilisco (✉)

Dipartimento SEFEMEQ and Centro V. Volterra, Facoltà di Economia,  
Università di Roma “Tor Vergata”, Via Columbia 2, 00133 Rome, Italy  
e-mail: gibilisco@volterra.uniroma2.it

T. Isola

Dipartimento di Matematica, Università di Roma “Tor Vergata”, Via della Ricerca Scientifica,  
00133 Rome, Italy  
e-mail: isola@mat.uniroma2.it

No such lower bound for the variance of pairs of random variables exists in classical probability. Schrödinger proved a stronger inequality involving covariance

$$\text{Var}_\rho(A) \cdot \text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2 \geq \frac{1}{4} |\text{Tr}(\rho[A, B])|^2.$$

Recently S. Luo and Q. Zhang proved a different kind of uncertainty principle (see [Luo and Zhang \(2004\)](#), Theorem 2), in the Schrödinger form, where the lower bound appears because the variables  $A, B$  do not commute with the state  $\rho$  (in contrast with the standard uncertainty principle where the bound depends on the commutator  $[A, B]$ ).

The inequality was conjectured by S. Luo himself and Z. Zhang in a previous paper ([Luo and Zhang 2004](#)). These authors suggest there that “the result may be interpreted as a quantification of certain aspect of the Wigner–Araki–Yanase theorem for quantum measurement, which states that observables not commuting with a conserved quantity cannot be measured exactly” (see [Wigner 1952; Araki and Yanase 1960; Ozawa 2002](#)). The inequality has been recently generalized in [Kosaki \(2005\)](#) and [Yanagi et al. \(2005\)](#). The final result is

$$\text{Var}_\rho(A) \cdot \text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2 \geq I_{\rho, \beta}(A)I_{\rho, \beta}(B) - |\text{Re}\{\text{Corr}_{\rho, \beta}(A, B)\}|^2$$

where  $I$  and  $\text{Corr}$  are given by the Wigner–Yanase–Dyson skew information (see Sect. 3 below).

The purpose of this paper is to put the above inequality in a more geometric form by means of quantum Fisher information (namely the monotone metrics classified by Petz). In this way the lower bound will appear as a simple function of the area spanned by the commutators  $i[A, \rho], i[B, \rho]$  in the tangent space to the state  $\rho$ , provided the state space is equipped with a suitable monotone metric (see Theorem 6). At this point it is natural to ask whether such an inequality holds for other quantum Fisher informations in the Wigner–Yanase–Dyson class (like the *RLD*-metric for example). The answer turns out to be negative and a general counterexample is given in Proposition 2.

In the final section we discuss some open problems related to the subject.

## 2 Schrödinger and Heisenberg uncertainty principles

Let  $M_n := M_n(\mathbb{C})$  (resp.  $M_{n,sa} := M_n(\mathbb{C})_{sa}$ ) be the set of all  $n \times n$  complex matrices (resp. all  $n \times n$  self-adjoint matrices). We shall denote general matrices by  $X, Y, \dots$  while letters  $A, B, \dots$  will be used for self-adjoint matrices. Let  $\mathcal{D}_n$  be the set of strictly positive elements of  $M_n$  while  $\mathcal{D}_n^1 \subset \mathcal{D}_n$  is the set of strictly positive density matrices namely

$$\mathcal{D}_n^1 = \{\rho \in M_n | \text{Tr}\rho = 1, \rho > 0\}.$$

**Proposition 1** *The correspondence*

$$M_n \times M_n \ni (X, Y) \rightarrow \langle X, Y \rangle := \text{Tr}(\rho X Y^*) - \text{Tr}(\rho X) \cdot \overline{\text{Tr}(\rho Y)}$$

is a positive sesquilinear form.

As usual commutators and anticommutators are defined as  $[X, Y] = XY - YX$ ,  $\{X, Y\} = XY + YX$ .

**Definition 1** Suppose that  $\rho \in \mathcal{D}_n^1$  is fixed. Define  $X_0 := X - \text{Tr}(\rho X)I$ .

**Definition 2** For  $A, B \in M_{n,sa}$  and  $\rho \in \mathcal{D}_n^1$  define covariance and variance as

$$\begin{aligned} \text{Cov}_\rho(A, B) &:= \langle A, B \rangle = \text{Tr}(\rho AB) - \text{Tr}(\rho A) \cdot \text{Tr}(\rho B) = \text{Tr}(\rho A_0 B_0) \\ \text{Var}_\rho(A) &:= \langle A, A \rangle = \text{Tr}(\rho A^2) - \text{Tr}(\rho A)^2 = \text{Tr}(\rho A_0^2). \end{aligned}$$

Note that for  $A, B \in M_{n,sa}$  and  $\rho \in \mathcal{D}_n^1$  one has

$$\text{Re}(\text{Tr}(\rho AB)) = \frac{1}{2}\text{Tr}(\rho\{A, B\}) \quad \text{Im}(\text{Tr}(\rho AB)) = \frac{1}{2i}\text{Tr}(\rho[A, B]).$$

Since  $\text{Cov}_\rho(A, B) = \overline{\text{Cov}_\rho(B, A)}$  then

$$2\text{Re}\{\text{Cov}_\rho(A, B)\} = \text{Cov}_\rho(A, B) + \text{Cov}_\rho(B, A).$$

As a consequence of Cauchy–Schwartz inequality one can derive the Schrödinger and Heisenberg uncertainty principles that are given in the following

**Theorem 1** (see Schrödinger 1930) For  $A, B \in M_{n,sa}$  and  $\rho \in \mathcal{D}_n^1$  one has

$$\text{Var}_\rho(A) \cdot \text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2 \geq \frac{1}{4}|\text{Tr}(\rho[A, B])|^2$$

that implies

$$\text{Var}_\rho(A) \cdot \text{Var}_\rho(B) \geq \frac{1}{4}|\text{Tr}(\rho[A, B])|^2.$$

**Definition 3** Set

$$\mathcal{S}_\rho(A, B) := \text{Var}_\rho(A) \cdot \text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2.$$

*Remark 1* With the above definition the Schrödinger uncertainty principle takes the form

$$\mathcal{S}_\rho(A, B) \geq \frac{1}{4}|\text{Tr}(\rho[A, B])|^2.$$

Let us try to see this situation in general.

**Definition 4** Let  $F : \mathcal{D}_n^1 \times M_{n,sa} \times M_{n,sa} \rightarrow \mathbb{R}$  be a function (denoted as  $F_\rho(A, B)$ ) such that

$$\mathcal{S}_\rho(A, B) \geq F_\rho(A, B).$$

Then we say that  $F$  is an Uncertainty Principle Function (shortly UPF).

Problem are there nontrivial UPF different from  $\frac{1}{4}|\text{Tr}(\rho[A, B])|^2$ ?

More specifically the Heisenberg uncertainty principle says that if  $A$  and  $B$  do not commute then the product  $\text{Var}_\rho(A) \cdot \text{Var}_\rho(B)$  cannot be arbitrarily small. Is the same true if  $[A, \rho], [B, \rho] \neq 0$ ? The answer is given by Theorems 3, 6 and shows that in this case the bound on  $\text{Var}_\rho(A) \cdot \text{Var}_\rho(B)$  depends on a certain “area” spanned by the commutators  $[A, \rho], [B, \rho]$ .

### 3 An inequality related to uncertainty principle

**Definition 5** For  $A, B \in M_{n,sa}$ ,  $\rho \in \mathcal{D}_n^1$  and  $\beta \in (0, 1)$  set

$$\text{Corr}_{\rho, \beta}(A, B) := \text{Tr}(\rho AB) - \text{Tr}(\rho^\beta A \rho^{1-\beta} B).$$

With direct calculation one can prove the following

#### Lemma 1

$$\begin{aligned} 2\text{Re}\{\text{Corr}_{\rho, \beta}(A, B)\} &= \text{Corr}_{\rho, \beta}(A, B) + \text{Corr}_{\rho, \beta}(B, A) = -\text{Tr}([\rho^\beta A] \cdot [\rho^{1-\beta} B]), \\ \text{Corr}_{\rho, \beta}(A, B) &= \text{Cov}_\rho(A, B) - \text{Tr}(\rho^\beta A_0 \rho^{1-\beta} B_0). \end{aligned}$$

**Definition 6** The Wigner–Yanase–Dyson information is defined as

$$I_{\rho, \beta}(A) := \text{Corr}_{\rho, \beta}(A, A) = -\frac{1}{2}\text{Tr}([\rho^\beta A] \cdot [\rho^{1-\beta} A]).$$

#### Definition 7

$$T_{\rho, \beta}(A, B) := I_{\rho, \beta}(A)I_{\rho, \beta}(B) - |\text{Re}\{\text{Corr}_{\rho, \beta}(A, B)\}|^2.$$

Note that  $T_{\rho, \beta} = T_{\rho, 1-\beta}$  so one can consider just  $\beta \in (0, \frac{1}{2}]$ .

In [Luo and Zhang \(2004\)](#) the following result has been proved

**Theorem 2**  $T_{\rho, \frac{1}{2}}(A, B)$  is an UPF.

The theorem had been conjectured in [Luo and Zhang \(2004\)](#). A generalization of Theorem 2 has been given in [Kosaki \(2005\)](#) and [Yanagi et al. \(2005\)](#).

**Theorem 3**  $T_{\rho, \beta}(A, B)$  is an UPF for any  $\beta \in (0, \frac{1}{2}]$ .

*Proof* We report here the proof of Yanagi et al. (2005) because it is needed in the sequel.

We have to prove that for any two self-adjoint operators  $A$  and  $B$ , any density operator  $\rho$  and any  $0 < \beta \leq \frac{1}{2}$ , we have

$$\begin{aligned} \text{Var}_\rho(A) \text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2 \\ \geq I_{\rho, \beta}(A) I_{\rho, \beta}(B) - |\text{Re}\{\text{Corr}_{\rho, \beta}(A, B)\}|^2. \end{aligned}$$

Let  $\{\varphi_i\}$  be a complete orthonormal base composed of eigenvectors of  $\rho$ , and  $\{\lambda_i\}$  the corresponding eigenvalues.

Set  $a_{ij} \equiv \langle A_0 \varphi_i | \varphi_j \rangle$  and  $b_{ij} \equiv \langle B_0 \varphi_i | \varphi_j \rangle$ .

Then we calculate

$$\begin{aligned} \text{Var}_\rho(A) &= \text{Tr}(\rho A_0^2) = \frac{1}{2} \sum_{i,j} (\lambda_i + \lambda_j) a_{ij} a_{ji}, \\ \text{Var}_\rho(B) &= \text{Tr}(\rho B_0^2) = \frac{1}{2} \sum_{i,j} (\lambda_i + \lambda_j) b_{ij} b_{ji}, \\ \text{Re}\{\text{Cov}_\rho(A, B)\} &= \text{Re}\{\text{Tr}(\rho A_0 B_0)\} = \frac{1}{2} \sum_{i,j} (\lambda_i + \lambda_j) \text{Re}\{a_{ij} b_{ji}\}, \\ I_{\rho, \beta}(A) &= \text{Var}_\rho(A) - \text{Tr}(\rho^\beta A_0 \rho^{1-\beta} A_0) \\ &= \frac{1}{2} \sum_{i,j} (\lambda_i + \lambda_j) a_{ij} a_{ji} - \sum_{i,j} \lambda_i^\beta \lambda_j^{1-\beta} a_{ij} a_{ji}, \\ I_{\rho, \beta}(B) &= \frac{1}{2} \sum_{i,j} (\lambda_i + \lambda_j) b_{ij} b_{ji} - \sum_{i,j} \lambda_i^\beta \lambda_j^{1-\beta} b_{ij} b_{ji}, \\ \text{Re}\{\text{Corr}_{\rho, \beta}(A, B)\} &= \text{Re}\{\text{Cov}_\rho(A, B)\} - \text{Re}\{\text{Tr}(\rho^\beta A_0 \rho^{1-\beta} B_0)\} \\ &= \frac{1}{2} \sum_{i,j} (\lambda_i + \lambda_j) \text{Re}\{a_{ij} b_{ji}\} - \sum_{i,j} \lambda_i^\beta \lambda_j^{1-\beta} \text{Re}\{a_{ij} b_{ji}\}. \end{aligned}$$

Set

$$\begin{aligned} \xi &:= \text{Var}_\rho(A) \text{Var}_\rho(B) - I_{\rho, \beta}(A) I_{\rho, \beta}(B) \\ &= \frac{1}{2} \sum_{i,j,k,l} \left\{ (\lambda_i + \lambda_j) \lambda_k^\beta \lambda_l^{1-\beta} + (\lambda_k + \lambda_l) \lambda_i^\beta \lambda_j^{1-\beta} - 2 \lambda_i^\beta \lambda_j^{1-\beta} \lambda_k^\beta \lambda_l^{1-\beta} \right\} a_{ij} a_{ji} b_{kl} b_{lk} \\ &= \frac{1}{4} \sum_{i,j,k,l} \left\{ (\lambda_i + \lambda_j) \lambda_k^\beta \lambda_l^{1-\beta} + (\lambda_k + \lambda_l) \lambda_i^\beta \lambda_j^{1-\beta} - 2 \lambda_i^\beta \lambda_j^{1-\beta} \lambda_k^\beta \lambda_l^{1-\beta} \right\} \\ &\quad \times \{a_{ij} a_{ji} b_{kl} b_{lk} + a_{kl} a_{lk} b_{ij} b_{ji}\}, \end{aligned}$$

$$\begin{aligned}\eta &:= |\operatorname{Re} \{\operatorname{Cov}_\rho(A, B)\}|^2 - |\operatorname{Re} \{\operatorname{Corr}_{\rho, \beta}(A, B)\}|^2 \\ &= \frac{1}{2} \sum_{i,j,k,l} \left\{ (\lambda_i + \lambda_j) \lambda_k^\beta \lambda_l^{1-\beta} + (\lambda_k + \lambda_l) \lambda_i^\beta \lambda_j^{1-\beta} - 2 \lambda_i^\beta \lambda_j^{1-\beta} \lambda_k^\beta \lambda_l^{1-\beta} \right\} \\ &\quad \times \operatorname{Re}\{a_{ij} b_{ji}\} \operatorname{Re}\{a_{kl} b_{lk}\}.\end{aligned}$$

In order to prove the theorem it is enough to show  $\xi - \eta \geq 0$ . Indeed

$$\begin{aligned}\xi - \eta &= \frac{1}{4} \sum_{i,j,k,l} \left\{ (\lambda_i + \lambda_j) \lambda_k^\beta \lambda_l^{1-\beta} + (\lambda_k + \lambda_l) \lambda_i^\beta \lambda_j^{1-\beta} - 2 \lambda_i^\beta \lambda_j^{1-\beta} \lambda_k^\beta \lambda_l^{1-\beta} \right\} \\ &\quad \cdot \left\{ |a_{ij}|^2 |b_{kl}|^2 + |a_{kl}|^2 |b_{ij}|^2 - 2 \operatorname{Re}\{a_{ij} b_{ji}\} \operatorname{Re}\{a_{kl} b_{lk}\} \right\}.\end{aligned}$$

Since

$$\begin{aligned}(\lambda_i + \lambda_j) \lambda_k^\beta \lambda_l^{1-\beta} + (\lambda_k + \lambda_l) \lambda_i^\beta \lambda_j^{1-\beta} - 2 \lambda_i^\beta \lambda_j^{1-\beta} \lambda_k^\beta \lambda_l^{1-\beta} \\ = (\lambda_i + \lambda_j - \lambda_i^\beta \lambda_j^{1-\beta}) \lambda_k^\beta \lambda_l^{1-\beta} + (\lambda_k + \lambda_l - \lambda_k^\beta \lambda_l^{1-\beta}) \lambda_i^\beta \lambda_j^{1-\beta} \geq 0, \\ |a_{ij}|^2 |b_{kl}|^2 + |a_{kl}|^2 |b_{ij}|^2 \geq 2 |a_{ij} b_{ji}| |a_{kl} b_{lk}| \geq 2 |\operatorname{Re} \{a_{ij} b_{ji}\} \operatorname{Re} \{a_{kl} b_{lk}\}|,\end{aligned}$$

we get the thesis.

*Remark 2* Note that Kosaki proved Theorem 3 by showing that  $T_{\rho, \beta}(A, B)$  is monotone increasing for  $\beta \in (0, \frac{1}{2}]$ . Moreover he was able to prove that  $S_\rho(A, B) = T_{\rho, \beta}(A, B)$  iff  $A_0, B_0$  are proportional.

#### 4 A counterexample

The inequality of Theorem 3 is not true for arbitrary values of  $\beta$  as it is proved in the following

**Proposition 2** *For any  $\beta \in [-1, 0)$  there are a state  $\rho$  and self-adjoint operators  $A$  and  $B$  s.t.*

$$\begin{aligned}\operatorname{Var}_\rho(A) \operatorname{Var}_\rho(B) - |\operatorname{Re} \{\operatorname{Cov}_\rho(A, B)\}|^2 &< I_{\rho, \beta}(A) I_{\rho, \beta}(B) \\ - |\operatorname{Re} \{\operatorname{Corr}_{\rho, \beta}(A, B)\}|^2.\end{aligned}$$

*Proof 2* Let  $t \in (0, \frac{1}{2})$  and

$$\rho = \begin{pmatrix} t & 0 & 0 \\ 0 & 1-2t & 0 \\ 0 & 0 & t \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Then, using the calculations performed for the proof of Theorem 3, we have

$$\begin{aligned}
\xi - \eta &= \text{Var}_\rho(A)\text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2 - I_{\rho,\beta}(A)I_{\rho,\beta}(B) \\
&\quad - |\text{Re}\{\text{Corr}_{\rho,\beta}(A, B)\}|^2 \\
&= \frac{1}{2} \sum_{i,j,k,l} \left\{ (\lambda_i + \lambda_j) \lambda_k^\beta \lambda_\ell^{1-\beta} + (\lambda_k + \lambda_\ell) \lambda_i^\beta \lambda_j^{1-\beta} - 2\lambda_i^\beta \lambda_j^{1-\beta} \lambda_k^\beta \lambda_\ell^{1-\beta} \right\} \\
&\quad \cdot \{a_{ij}a_{ji}b_{kl}b_{lk} - \text{Re}(a_{ij}b_{ji})\text{Re}(a_{kl}b_{lk})\} \\
&= \frac{1}{2} (2\lambda_1 + 2\lambda_2 - \lambda_1^\beta \lambda_2^{1-\beta} - \lambda_2^\beta \lambda_1^{1-\beta})(\lambda_2^\beta \lambda_3^{1-\beta} + \lambda_3^\beta \lambda_2^{1-\beta}) \\
&\quad + \frac{1}{2} (2\lambda_2 + 2\lambda_3 - \lambda_2^\beta \lambda_3^{1-\beta} - \lambda_3^\beta \lambda_2^{1-\beta})(\lambda_1^\beta \lambda_2^{1-\beta} + \lambda_2^\beta \lambda_1^{1-\beta}) \\
&= \{2(1-t) - t^\beta(1-2t)^{1-\beta} - (1-2t)^\beta t^{1-\beta}\} \{t^\beta(1-2t)^{1-\beta} + (1-2t)^\beta t^{1-\beta}\}.
\end{aligned}$$

Let  $\beta \in [-1, 0]$ . Since  $t^\beta(1-2t)^{1-\beta} \rightarrow \infty$  if  $t \rightarrow 0^+$ , there exists a  $t_0 = t_0(\beta) \in (0, 1)$  for which  $\xi - \eta < 0$ . This ends the proof.

*Remark 3* For  $\beta \in [-1, 0)$  the inequality

$$\begin{aligned}
\text{Var}_\rho(A)\text{Var}_\rho(B) - |\text{Re}\{\text{Cov}_\rho(A, B)\}|^2 &< I_{\rho,\beta}(A)I_{\rho,\beta}(B) \\
- |\text{Re}\{\text{Corr}_{\rho,\beta}(A, B)\}|^2
\end{aligned}$$

is not true in general as one can see by choosing

$$t \in (0, 1), \quad \rho = \begin{pmatrix} t & 0 \\ 0 & 1-t \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

In the next Sections we try to give a more geometric form to Theorem 3 and Proposition 2.

## 5 Quantum Fisher informations

In what follows if  $\mathcal{N}$  is a differential manifold we denote by  $T_\rho \mathcal{N}$  the tangent space to  $\mathcal{N}$  at the point  $\rho \in \mathcal{N}$ .

In the commutative case a Markov morphism is a stochastic map  $T : \mathbb{R}^n \rightarrow \mathbb{R}^k$ . In the noncommutative case a Markov morphism is a completely positive and trace preserving operator  $T : M_n \rightarrow M_k$ . Let

$$\mathcal{P}_n := \{\rho \in \mathbb{R}^n | \rho_i > 0\}, \quad \mathcal{P}_n^1 := \left\{ \rho \in \mathbb{R}^n | \sum \rho_i = 1, \rho_i > 0 \right\}.$$

In the commutative case a monotone metric is a family of riemannian metrics  $g = \{g^n\}$  on  $\{\mathcal{P}_n^1\}$ ,  $n \in \mathbb{N}$ , such that

$$g_{T(\rho)}^m(TX, TX) \leq g_\rho^n(X, X)$$

holds for every Markov morphism  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and all  $\rho \in \mathcal{P}_n^1$  and  $X \in T_\rho \mathcal{P}_n^1$ .

It is not difficult to see that there exists a natural identification of  $T_\rho \mathcal{D}_n^1$  with the space of self-adjoint traceless matrices, namely

$$T_\rho \mathcal{D}_n^1 = \{A \in M_n | A = A^*, \text{ Tr}(A) = 0\}.$$

In perfect analogy with the commutative case, a monotone metric in the non-commutative case is a family of riemannian metrics  $g = \{g^n\}$  on  $\{\mathcal{D}_n^1\}$ ,  $n \in \mathbb{N}$ , such that

$$g_{T(\rho)}^m(TX, TX) \leq g_\rho^n(X, X)$$

holds for every Markov morphism  $T : M_n \rightarrow M_m$  and all  $\rho \in \mathcal{D}_n^1$  and  $X \in T_\rho \mathcal{D}_n^1$ .

Let us recall that a function  $f : (0, \infty) \rightarrow \mathbb{R}$  is said operator monotone if, for any  $n \in N$ , any  $A, B \in M_n$  such that  $0 \leq A \leq B$ , the inequalities  $0 \leq f(A) \leq f(B)$  hold. An operator monotone function is said symmetric if  $f(x) := xf(x^{-1})$ . With such operator monotone functions  $f$  one associates the so-called Chentsov–Morotzova functions

$$c_f(x, y) := \frac{1}{yf(xy^{-1})} \quad \text{for } x, y > 0.$$

Define  $L_\rho(A) := \rho A$ , and  $R_\rho(A) := A\rho$ . Since  $L_\rho$  and  $R_\rho$  commute we may define  $c_f(L_\rho, R_\rho)$ . Now we can state the fundamental theorems about monotone metrics. In what follows uniqueness and classification are stated up to scalars (see Petz 1996).

**Theorem 4** (Chentsov 1982) *There exists a unique monotone metric on  $\mathcal{P}_n^1$  given by the Fisher information.*

**Theorem 5** (Petz 1996) *There exists a bijective correspondence between symmetric monotone metrics on  $\mathcal{D}_n^1$  and symmetric operator monotone functions. This correspondence is given by the formula*

$$g_f(A, B) := g_{f, \rho}(A, B) := \text{Tr}(A \cdot c_f(L_\rho, R_\rho)(B)).$$

Because of these two theorems we shall use the terms “Monotone metrics” and “Quantum Fisher informations” (shortly QFI) with the same meaning.

Note that usually monotone metrics are normalized so that if  $[A, \rho] = 0$  then  $g_{f, \rho}(A, A) = \text{Tr}(\rho^{-1} A^2)$ , that is equivalent to ask  $f(1) = 1$ .

Examples of monotone metrics are given by the following list (see Hasegawa and Petz 1997; Gibilisco and Isola 2004).

Let

$$\begin{aligned} f_\beta(x) &:= \beta(1-\beta) \frac{(x-1)^2}{(x^\beta - 1)(x^{1-\beta} - 1)}, \quad \beta \in [-1, \frac{1}{2}] \setminus \{0\}, \\ f_0(x) &:= \frac{x-1}{\log x}, \\ h_\gamma(x) &:= \left(\frac{1+x^\gamma}{2}\right)^{\frac{1}{\gamma}} \quad \gamma \in [\frac{1}{2}, 1]. \end{aligned}$$

Note that  $f_0 = \lim_{\beta \rightarrow 0} f_\beta$ .

The metrics associated with the functions  $f_\beta$  are equivalent to the metrics induced by noncommutative  $\alpha$ -divergences where  $\beta = \frac{1-\alpha}{2}$  (see Hasegawa and Petz 1997).

The *RLD*-metric is the QFI associated to  $f_{-1}$ .

The *BKM*-metric is the QFI associated to  $f_0$ .

The *WY*-metric is the QFI associated to  $f_{\frac{1}{2}} = h_{\frac{1}{2}}$ .

The *SLD*-metric (or Bures-Uhlmann metric) is the QFI associated to  $h_1$ .

The two parametric families  $f_\beta, h_\gamma$  give us a continuum of operator monotone functions from the smallest  $f_{-1}(x) = \frac{2x}{x+1}$  to the greatest  $h_1 = \frac{1+x}{2}$ .

For a symmetric operator monotone function  $\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = f(0) := \lim_{x \rightarrow 0} f(x)$ . Note that

$$\begin{aligned} f_\beta(0) &= 0 \quad \beta \in [-1, 0], \\ f_\beta(0) &= \beta(1-\beta) \neq 0 \quad \beta \in (0, \frac{1}{2}], \\ h_\gamma(0) &= \left(\frac{1}{2}\right)^{\frac{1}{\gamma}} \neq 0 \quad \gamma \in [\frac{1}{2}, 1]. \end{aligned}$$

The condition  $f(0) \neq 0$  is relevant because it is a necessary and sufficient condition for the existence of the so-called radial extension of a monotone metric to pure states (see Petz and Sudar 1996).

## 6 A geometric look at the inequality

Let  $V$  be a finite dimensional real vector space with a scalar product  $g(\cdot, \cdot)$ . We define, for  $v, w \in V$ ,

$$\text{Area}_g(v, w) := \sqrt{g(v, v) \cdot g(w, w) - |g(v, w)|^2}.$$

In the euclidean plane  $\text{Area}_g(v, w)$  is the area of the parallelogramme spanned by  $v$  and  $w$ .

Define  $A_\rho := i[\rho, A]$ . Since  $A_\rho$  is traceless and selfadjoint, then  $A_\rho \in T_\rho \mathcal{D}_n^1$ .

**Proposition 3** *For the QFI associated to  $f_\beta$  one has*

$$g_\beta(A_\rho, B_\rho) := g_{f_\beta}(A_\rho, B_\rho) = -\frac{1}{\beta(1-\beta)} \text{Tr}([\rho^\beta, A] \cdot [\rho^{1-\beta}, B]) \quad \beta \in \left[-1, \frac{1}{2}\right] \setminus \{0\}.$$

One can find a proof in Hasegawa and Petz (1997), Gibilisco and Isola (2004). Because of the above proposition  $g_\beta$  is known as the WYD( $\beta$ ) monotone metric.

If  $f$  is an operator monotone function we denote by  $\text{Area}_f$  the area functional associated to the monotone metric  $g_f$ . One has

### Theorem 6

$$\mathcal{T}_{\rho, \beta}(A, B) = \frac{(\beta(1-\beta))^2}{4} (\text{Area}_{f_\beta}(i[\rho, A], i[\rho, B]))^2 \quad \forall \beta \in \left[-1, \frac{1}{2}\right] \setminus \{0\}.$$

*Proof 3* One has from Lemma 1 and Proposition 3

$$\begin{aligned} \mathcal{T}_{\rho, \beta}(A, B) &= I_{\rho, \beta}(A)I_{\rho, \beta}(B) - |\text{Re} \{ \text{Corr}_{\rho, \beta}(A, B) \}|^2 \\ &= \left( -\frac{1}{2} \text{Tr}([\rho^\beta, A] \cdot [\rho^{1-\beta}, A]) \right) \cdot \left( -\frac{1}{2} \text{Tr}([\rho^\beta, B] \cdot [\rho^{1-\beta}, B]) \right) \\ &\quad - \frac{1}{4} |\text{Tr}([\rho^\beta, A] \cdot [\rho^{1-\beta}, B])|^2 \\ &= \frac{(\beta(1-\beta))^2}{4} (g_\beta(A_\rho, A_\rho) \cdot g_\beta(B_\rho, B_\rho) - |g_\beta(A_\rho, B_\rho)|^2) \\ &= \frac{(\beta(1-\beta))^2}{4} (\text{Area}_{f_\beta}(i[\rho, A], i[\rho, B]))^2. \end{aligned}$$

## 7 Relation with curvature

The appearance of the area of a Riemannian metric in Theorem 6 (and therefore in Theorem 3) suggests a link between the uncertainty principle and the notion of curvature. In this section we make some considerations of this point. To make the paper self-contained we recall some notions of differential geometry.

For an affine (linear) connection  $\nabla$  on a manifold  $\mathcal{M}$  the curvature is defined as (see Kobayashi and Nomizu 1963, p. 133)

$$R(X, Y)Z := [\nabla_X, \nabla_Y]Z - \nabla_{[X, Y]}Z.$$

Suppose that  $g(\cdot, \cdot)$  is a Riemannian metric on  $\mathcal{M}$  and  $\nabla$  is the associated Levi-Civita connection. The Riemannian curvature tensor is defined as (see Kobayashi and Nomizu 1963, p. 201)

$$R(X, Y, Z, W) := g(R(Z, W)Y, X)$$

where  $X, Y, Z, W$  are vector fields.

Now let  $\rho \in \mathcal{M}$  and suppose that we have a 2-dimensional subspace  $\sigma \subset T_\rho \mathcal{M}$ . Then  $\sigma$  determines, with the use of the exponential map  $\exp$ , a 2-dimensional embedded surface  $\mathcal{N} := \exp_\rho(B_\eta(0_\rho) \cap \sigma)$  formed by the geodesic segments of length  $< \eta$  which start tangentially to  $\sigma$ . If  $K(\sigma)$  denotes the Gaussian curvature of  $\mathcal{N}$  one has the following

**Proposition 4** (see Klingenberg 1982, p. 99–100) *If  $A, B$  is a basis for the plane  $\sigma$  then*

$$K(A, B) := K(\sigma) = \frac{R(A, B, A, B)}{g(A, A)g(B, B) - |g(A, B)|^2} = \frac{g(R(A, B)B, A)}{\text{Area}_g(A, B)^2}.$$

When we want to emphasize the dependence of  $R$  and  $K$  from the Riemannian metric  $g$  we write  $R_g$  and  $K_g$ .

If  $f$  is an operator monotone function we denote by  $R_f$  the Riemannian curvature tensor and by  $K_f$  the sectional curvature.

Note that, if  $\beta = \frac{1}{2}$ , then  $K_{f_{\frac{1}{2}}}(\sigma) = \text{costant} = \frac{1}{4}$  (see Gibilisco and Isola 2003), so the inequality of Theorem 2

$$\mathcal{S}_\rho(A, B) \geq I_{\rho, \frac{1}{2}}(A)I_{\rho, \frac{1}{2}}(B) - |\text{Re} \left\{ \text{Corr}_{\rho, \frac{1}{2}}(A, B) \right\}|^2$$

takes the form

$$\mathcal{S}_\rho(A, B) \geq \frac{1}{16}R_{f_{\frac{1}{2}}}(A_\rho, B_\rho, A_\rho, B_\rho).$$

In general from bounds on sectional curvature  $K_{f_\beta}(\sigma)$  one would be able to deduce inequalities of the same type for the Riemann curvature tensor (see Gibilisco and Isola (2005) for ideas about this kind of bounds).

## 8 Conclusions and open problems

We can summarize Theorem 3, Proposition 2 and Theorem 6 into the following.

**Theorem 7**

$$\begin{aligned} \mathcal{S}_\rho(A, B) &\geq \frac{(\beta(1-\beta))^2}{4} (\text{Area}_{f_\beta}(i[\rho, A], i[\rho, B]))^2 \\ &\Updownarrow \\ \beta &\in \left[ 0, \frac{1}{2} \right] \end{aligned}$$

We have an inequality that is true only for some elements  $f_\beta$  of the class of the Wigner–Yanase–Dyson monotone metrics. For example it is true for the *WY-*

metric ( $\beta = \frac{1}{2}$ ) and is false for the *RLD*-metric ( $\beta = -1$ ). In this “iff” form the above inequality seems a result that cannot be further generalized.

### Problem 1

Maybe one should still seek a different generalization of Theorem 7. Since

$$\beta \in [-1, 0] \implies f_\beta(0) = 0 \quad \& \quad \beta \in (0, \frac{1}{2}] \implies f_\beta(0) = \beta(1 - \beta)$$

one can state Theorem 7 (that is Theorem 3) in a different way

### Theorem 8

$$\mathcal{S}_\rho(A, B) \geq \frac{f_\beta(0)^2}{4} (\text{Area}_{f_\beta}(i[A, \rho], i[B, \rho]))^2 \quad \forall \beta \in [-1, \frac{1}{2}]$$

Question: characterize the family of operator monotone functions  $f$  for which is true the inequality

$$\mathcal{S}_\rho(A, B) \geq \frac{f(0)^2}{4} (\text{Area}_f(i[A, \rho], i[B, \rho]))^2.$$

Of course the above inequality is trivially true when  $f(0) = 0$  while it is a non-trivial inequality for those operator monotone functions such that  $f(0) > 0$ . Note that the question is non-trivial, for example, for the *SLD*-metric for which  $h_1(0) = \frac{1}{2}$ .

### Problem 2

For  $f$  operator monotone define

$$G(f) := \frac{f(0)^2}{4} (\text{Area}_f(i[A, \rho], i[B, \rho]))^2.$$

In Kosaki (2005) the proof of Theorem 3 (see p. 640) is obtained by the following result

$$f_\beta \leq f_{\tilde{\beta}} \implies G(f_\beta) \leq G(f_{\tilde{\beta}}) \quad \beta \in (0, \frac{1}{2}]$$

Is this inequality true for other families of operator monotone functions?

### Problem 3

The following question has been posed at p. 642 in Kosaki (2005). Covariance and *WYD* information make perfect sense in infinite dimension (see Connes and Størmer 1978, Kosaki 1982), namely in a general von Neumann algebra setting. Is the inequality of Theorem 3 still true in this general setting?

**Acknowledgements** The present results have been presented at the second International Conference “Information Geometry and its Applications” held at the University of Tokio, December 12–16, 2005. It is a pleasure to thank all the organizers.

## References

- Araki, H., Yanase, M.M. (1960). Measurement of quantum mechanical operators. *Physical Review*, 120(2), 622–626.
- Connes, A., Størmer, E. (1978). Homogeneity of the state space of factors of type  $III_1$ . *Journal of Functional Analysis*, 28, 187–196.
- Gibilisco, P., Isola, T. (2003). Wigner-Yanase information on quantum state space: the geometric approach. *Journal of Mathematical Physics*, 44(9), 3752–3762.
- Gibilisco, P., Isola, T. (2004). On the characterisation of paired monotone metrics. *Annals of the Institute of Statistical Mathematics*, 56(2), 369–381.
- Gibilisco, P., Isola, T. (2005). On the monotonicity of scalar curvature in classical and quantum information geometry. *Journal of Mathematical Physics*, 46(2), 023501–14.
- Gibilisco, P., Isola, T. (2006). Some open problems in Information Geometry. In: *Proceedings 26th Conference: QP and IDA—Levico (Trento), February 20–26, 2005* (to appear)
- Hasegawa, H., Petz, D. (1997). Non-commutative extension of the information geometry II. In: O. Hirota et al. (Eds.), *Quantum Communications, Computing and Measurement* (pp. 109–118). New York: Plenum.
- Klingenberg, W. (1982). *Riemannian Geometry*. Berlin: Walter de Gruyter & Co.
- Kobayashi, S., Nomizu, K. (1963). *Foundations of differential geometry*, Vol. I. New York: John Wiley.
- Kosaki, H. (1982). Interpolation theory and the Wigner–Yanase–Dyson–Lieb concavity. *Communications in Mathematical Physics*, 87, 315–329.
- Kosaki, H. (2005). Matrix trace inequalities related to uncertainty principle. *International Journal of Mathematics*, 6, 629–645.
- Luo, S., Zhang, Q. (2004). On skew information. *IEEE Transactions on Information Theory*, 50(8), 1778–1782.
- Luo, S., Zhang, Z. (2004). An informational characterization of Schrödinger’s uncertainty relations. *Journal of Statistical Physics*, 114(5–6), 1557–1576.
- Ozawa, M. (2002). Conservation laws, uncertainty relations and quantum limits of measurement. *Physical Review Letters*, 88(5), 050402–4.
- Petz, D. (1996). Monotone metrics on matrix spaces, *Linear Algebra and Applications*, 244, 81–96.
- Petz, D., Sudar, C. (1996). Geometries of quantum states, *Journal of Mathematical Physics*, 37(6), 2662–2673.
- Schrödinger, E. (1930). About Heisenberg uncertainty relation (original annotation by A. Angelow and M.C. Batoni), *Bulgarian Journal of Physics*, 26(5–6), 193–203 (2000), 1999. Translation of *Proceedings Prussian Academy of Sciences, Physical and Mathematical Section*, 19 (1930), 296–303.
- Wigner, E. P. (1952). Die Messung quantenmechanischer Operatoren. *Zeitschrift für Physik*, 133, 101–108.
- Yanagi, K., Furuichi, S., Kuriyama, K. (2005). A generalized skew information and uncertainty relation. *IEEE Transactions on Information Theory*, 51(12), 4401–4404.